



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LOCAL RECORDS OF ARACHNIDS PERSONALLY COLLECTED IN KANSAS AND TEXAS.*

By E. S. TUCKER, Bureau of Entomology, U. S. Department of Agriculture.

A NUMBER of Arachnids collected by myself in two localities, one being Lawrence, Kan., and the other Plano, Tex., are herewith listed for local comparison. No special effort was made for collecting these creatures, and only such specimens as happened to be noticed at odd times were taken. The collection of the Kansas specimens preceded that from Plano, Tex., which were obtained during the season of 1907.

With further regard to the Kansas spiders, a revision of the specific names as given by Mr. T. H. Scheffer is made to conform so far as possible with the "Catalogue of Nearctic Spiders," by Mr. Nathan Banks (Bull. 72, U. S. Nat. Mus.). The list includes the following species:

Family CLUBIONIDÆ.

Trachelas tranquilla Htz. September, from beehive infested with wax-worms and moths (*Galleria mellonella* L.).

Anyphæna gracilis Htz. September.

Family AGELENIDÆ.

Agelena nævia Walck. August and September.

Family THERIDIIDÆ.

Theridium tepidariorum Koch. September.

Teutana triangulosa Walck. July, August and September.

Family TETRAGNATHIDÆ.

Tetragnatha extensa L. May and June.

Family EPEIRIDÆ.

Leucauge hortorum Htz. June, July and August.

Cyclosa conica Pal. July.

Mangora gibberosa Htz. August and September.

Epeira gigas Leach. October, web spanning window.

domiciliorum Htz. August and September.

globosa Keys. July.

ocellata Clerck. April; and July, August and September, occurring near electric lights on bridge.

trivittata Keys. September.

verrucosa Htz. August.

Plectana stellata Htz. April.

Argiope trifasciata Forsk. July, September and October.

* Published by permission of the chief of the Bureau of Entomology, U. S. Department of Agriculture.

Acrosoma gracilis Walck. July and August.
mitrata Htz. August.
spinea Htz. July.

Family THOMISIDÆ.

Xysticus gulosus Keys. September.
triguttatus Keys. June.
Runcinia aleatoria Htz. July and August.
Misumena vatia Clerck. September. New to Kansas list.
Misumessus asperatus Htz. April, May, July, August and September.
Tibellus oblongus Walck. April.

Family PISAURIDÆ.

Dolomedes mira Walck. August and September. (Not known to Banks.)

Family LYCOSIDÆ.

Lycosa avida Walck. September.
helluo Walck. April, young; July, female carrying cocoon; September, from beehive infested with waxworms and moths (*Galleria mellonella* L.); and October.

Family OXYOPIDÆ.

Oxyopes salticus Htz. July.

Family ATTIDÆ.

Phidippus audax Htz. April, July, September and October. Also, in October, dragging away a moth of *Galleria mellonella* L., on ceiling of room.
Dendryphantes octavus Htz. October.
Hycia pickei Peck. October.
Marpissa undata DeG. May, September.

All of the following species from Plano, Tex., were determined by Mr. Nathan Banks:

ORDER SCORPIONIDA: Family SCORPIONIDÆ.

Centrurus vittatus Say. May and July, under litter and in house.

ORDER PSEUDOSCORPIONES: Family CHERNETIDÆ.

Chelifer cancroides L. July and August.

ORDER PHALANGIIDEA: Family PHALANGIIDÆ.

Mesosoma niger Say. Summer and fall months.

ORDER ARANEIDA.

(Spiders collected July to November.)

Family SCYTODIDÆ.

Loxosceles rufescens Duf.

Family THERIDIIDÆ.

Lathrodictes mactans Fab.

Family TETRAGNATHIDÆ.

Tetragnatha grallator Htz.

Family EPEIRIDÆ.

Epeira foliata Koch.

gigas Leach.

prompta Htz.

Argiope aurantia Lucas.

Family THOMISIDÆ.

Runcinia aleatoria Htz.

Family LYCOSIDÆ.

Lycosa carolinensis Walck.

rabida Walck.

Family OXYOPIDÆ.

Peucetia viridans Htz.

Family ATTIDÆ.

Phidippus audax Htz.

Ashtabula, sp. (Young.) May.